Reg. No. :

## Question Paper Code : 21522

## B.E./B.Tech. DEGREE EXAMINATION, MAY/JUNE 2013.

Third Semester

**Civil Engineering** 

## 

(Common to all branches)

(Regulation 2008/2010)

Time : Three hours

Maximum: 100 marks

Answer ALL questions.

$$PART A - (10 \times 2 = 20 \text{ marks})$$

1. State the Dirichlet's conditions for Fourier series.

2. What is meant by Harmonic Analysis?

3. Find the Fourier Sine Transform of  $e^{-3x}$ .

4. If 
$$\mathfrak{Z}_{\mathfrak{l}}\{f(x)\}=F(s)$$
, prove that  $\mathfrak{Z}_{\mathfrak{l}}\{f(ax)\}=\frac{1}{a}\cdot F\left(\frac{s}{a}\right)$ 

5. Form the PDE from 
$$(x-a)^2 + (y-b)^2 + z^2 = r^2$$
.

6. Find the complete integral of p + q = pq.

7. In the one dimensional heat equation  $u_t = c^2 \cdot u_{xx}$ , what is  $c^2$ ?

8. Write down the two dimensional heat equation both in transient and steady states.

9. Find Z(n).

10. Obtain 
$$Z^{-1} \left[ \frac{z}{(z+1)(z+2)} \right]$$
.  
PART B — (5 × 16 = 80 marks)

11. (a) (i) Find the Fourier series of  $x^2$  in  $(-\pi, \pi)$  and hence deduce that  $\frac{1}{1^4} + \frac{1}{2^4} + \frac{1}{3^4} + \dots \frac{\pi^4}{90}.$ (8)

(ii) Obtain the Fourier cosine series of  $f(x) = \begin{cases} kx, & 0 < x < \frac{l}{2} \\ k(l-x), & \frac{l}{2} < x < l \end{cases}$  (8)

- (b) (i) Find the complex form of Fourier series of  $\cos ax$  in  $(-\pi, \pi)$ , where "a" is not an integer. (8)
  - (ii) Obtain the Fourier cosine series of  $(x-1)^2$ , 0 < x < 1 and hence show

that 
$$\frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \dots = \frac{\pi^2}{6}$$
. (8)

12. (a) (i) Find the Fourier transform of  $f(x) = \begin{cases} 1, |x| < a \\ 0, |x| > a \end{cases}$  and hence find

$$\int_{0} \frac{\sin x}{x} dx.$$
 (8)

(ii) Verify the convolution theorem under Fourier Transform, for  $f(x)=g(x)=e^{-x^2}$ . (8)

Or

(b)

1

15.

(i) Obtain the Fourier Transform of  $e^{-x^2/2}$ . (8)

(ii) Evaluate 
$$\int_{0}^{\infty} \frac{dx}{(x^2 + a^2)^2}$$
 using Parseval's identity. (8)

3. (a) (i) Solve: 
$$x(y^2 - z^2)p + y(z^2 - x^2)q = z(x^2 - y^2)$$
. (8)  
(ii) Solve:  $(D^2 + DD' - 6D'^2)z - y \cos x$  (8)

(ii) Solve: 
$$(D^2 + DD' - 6D'^2)z = y \cos x$$
. (8)

(b) (i) Solve: 
$$z = px + qy + \sqrt{p^2 + q^2 + 1}$$
 (8)

(ii) Solve: 
$$(D^3 - 7DD'^2 - 6D'^3)z = \sin(2x + y).$$
 (8)

14. (a) A tightly stretched string between the fixed end points x = 0 and x = l is initially at rest in its equilibrium position. If each of its points is given a velocity kx(l-x), find the displacement y(x,t) of the string.

Or

(b) An infinitely long rectangular plate is of width 10 cm. The temperature along the short edge y = 0 is given by

 $u = \begin{cases} 20x, & 0 < x < 5\\ 20(10-x), 5 < x < 10 \end{cases}$  If all the other edges are kept at zero

temperature, find the steady state temperature at any point on it.

(a) (i) Find 
$$Z(\cos n \theta)$$
 and hence deduce  $Z\left(\cos \frac{n \pi}{2}\right)$ . (8)

(ii) Using Z-transform solve:  $y_{n+2} - 3y_{n+1} - 10y_n = 0$ ,  $y_0 = 1$  and  $y_1 = 0$ . (8)

Or

(b) (i) State and prove the second shifting property of 
$$Z$$
-transform. (6)  
(ii) Using convolution theorem, find  $Z^{-1}\left[\frac{z^2}{(z-a)(z-b)}\right]$ . (10)