

PART B — (5 × 16 = 80 marks)

11. (a) (i) What is the angle between the surfaces $x^2 + y^2 + z^2 = 9$ and $x^2 + y^2 - z - 3 = 0$ at the point $(2, -1, 2)$? (8)

(ii) Using Stoke's theorem, evaluate $\int_C xy dx + xy^2 dy$ taking C is a square bounded by the lines $x = 1, x = -1, y = 1, y = -1$. (8)

Or

(b) (i) Show that $\vec{F} = (z^2 + 2x + 3y)\vec{i} + (3x + 2y + z)\vec{j} + (y + 2xz)\vec{k}$ is irrotational and hence find the corresponding potential function ϕ . (8)

(ii) Using Gauss Divergence theorem, evaluate $\iiint \text{curl } \vec{F} \cdot \vec{n} dS$ where $\vec{F} = 4xz\vec{i} - y^2\vec{j} + yz\vec{k}$ over the cube bounded by $x = 0, x = 1, y = 0, y = 1, z = 0, z = 1$. (8)

12. (a) (i) Solve $(D^2 - 4D + 3)y = \sin 3x + x^2$. (8)

(ii) Using method of variation of parameters, solve $(D^2 + 1)y = \sec x$. (8)

Or

(b) (i) Solve $\frac{dx}{dt} + \frac{dy}{dt} + 5x + 7y = 2e^{2t}, 2\frac{dx}{dt} + 3\frac{dy}{dt} + x + y = e^t$. (8)

(ii) Solve $(x^2 D^2 - 3xD + 4)y = x^2$. (8)

13. (a) (i) Find the Laplace transform of $t^2 e^t \sin 4t$. (8)

(ii) Using Convolution theorem, find the inverse Laplace transform of $\frac{1}{(s+1)(s^2+1)}$. (8)

Or

(b) (i) Find the Laplace transform of $f(t) = \begin{cases} t: 0 < t < 1 \\ 0: 1 < t < 2 \end{cases}$ and $f(t+2) = f(t)$. (8)

(ii) Solve $y'' + 4y' + 8y = 1$, $y(0) = 0$, $y'(0) = 1$ by using Laplace transforms. (8)

14. (a) (i) Find the analytic function $f(z) = u + iv$ if $u = e^{x^2 - y^2} \cos 2xy$. Hence find v . (8)

(ii) Find the image in the w plane of the region of the z plane bounded by the straight lines $x = 1$, $y = 1$, $x + y = 1$ under the transformation $w = z^2$. (8)

Or

(b) (i) If $w = u(x, y) + iv(x, y)$ is an analytic function, the curves of the family $u(x, y) = a$ and the curves of the family $v(x, y) = b$ cut orthogonally, where a and b are constants. (8)

(ii) Find the bilinear transformation which maps the points $z = 0$, $z = 1$, $z = \infty$ in to the points $w = i$, $w = 1$, $w = -i$ respectively. (8)

15. (a) (i) Use Cauchy's integral formula to evaluate $\int_C \frac{z+1}{z^3 - 2z^2} dz$, where C is the circle $C: |z - 2 - i| = 2$. (8)

(ii) Find the Laurent's series of $f(z) = \frac{z}{(z-1)(z-2)}$ valid in the region $|z+2| < 3$ and $3 < |z+2| < 4$. (8)

Or

(b) Evaluate $\int_{-\infty}^{\infty} \frac{x^2}{(x^2 + a^2)(x^2 + b^2)} dx$, $a > b > 0$, by using contour integration. (16)

